

RELIABILITY

```
/VARIABLES=Erstrater_A_Ges Zweitater_A_Ges Drittrater_A_Ges  
/SCALE('ALL VARIABLES') ALL  
/MODEL=ALPHA  
/ICC=MODEL(MIXED) TYPE(CONSISTENCY) CIN=95 TESTVAL=0.
```

## Reliability

### Scale: ALL VARIABLES

#### Case Processing Summary

		N	%
Cases	Valid	31	100,0
	Excluded <sup>a</sup>	0	,0
	Total	31	100,0

a. Listwise deletion based on all variables in the procedure.

#### Reliability Statistics

Cronbach's Alpha	N of Items
,707	3

#### Intraclass Correlation Coefficient

	Intraclass Correlation <sup>b</sup>	95% Confidence Interval		F Test with True Value 0			
		Lower Bound	Upper Bound	Value	df1	df2	Sig
Single Measures	,445 <sup>a</sup>	,226	,652	3,410	30	60	,000
Average Measures	,707 <sup>c</sup>	,468	,849	3,410	30	60	,000

Two-way mixed effects model where people effects are random and measures effects are fixed.

- a. The estimator is the same, whether the interaction effect is present or not.
- b. Type C intraclass correlation coefficients using a consistency definition. The between-measure variance is excluded from the denominator variance.
- c. This estimate is computed assuming the interaction effect is absent, because it is not estimable otherwise.

RELIABILITY

```
/VARIABLES=Erstrater_A_Ges zweitater_A_Ges  
/SCALE('ALL VARIABLES') ALL  
/MODEL=ALPHA  
/ICC=MODEL(MIXED) TYPE(CONSISTENCY) CIN=95 TESTVAL=0.
```

## Reliability

### Scale: ALL VARIABLES

**Case Processing Summary**

		N	%
Cases	Valid	31	100,0
	Excluded <sup>a</sup>	0	,0
	Total	31	100,0

a. Listwise deletion based on all variables in the procedure.

**Reliability Statistics**

Cronbach's Alpha	N of Items
,765	2

**Intraclass Correlation Coefficient**

	Intraclass Correlation <sup>b</sup>	95% Confidence Interval		F Test with True Value 0			
		Lower Bound	Upper Bound	Value	df1	df2	Sig
Single Measures	,619 <sup>a</sup>	,344	,796	4,246	30	30	,000
Average Measures	,765 <sup>c</sup>	,512	,886	4,246	30	30	,000

Two-way mixed effects model where people effects are random and measures effects are fixed.

- a. The estimator is the same, whether the interaction effect is present or not.
- b. Type C intraclass correlation coefficients using a consistency definition. The between-measure variance is excluded from the denominator variance.
- c. This estimate is computed assuming the interaction effect is absent, because it is not estimable otherwise.

RELIABILITY

```

/VARIABLES=Erstrater_A_Ges Drittrater_A_Ges
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/ICC=MODEL(MIXED) TYPE(CONSISTENCY) CIN=95 TESTVAL=0.
    
```

**Reliability**

**Scale: ALL VARIABLES**

**Case Processing Summary**

		N	%
Cases	Valid	31	100,0
	Excluded <sup>a</sup>	0	,0
	Total	31	100,0

a. Listwise deletion based on all variables in the procedure.

### Reliability Statistics

Cronbach's Alpha	N of Items
,463	2

### Intraclass Correlation Coefficient

	Intraclass Correlation <sup>b</sup>	95% Confidence Interval		F Test with True Value 0			
		Lower Bound	Upper Bound	Value	df1	df2	Sig
Single Measures	,301 <sup>a</sup>	-,054	,589	1,863	30	30	,047
Average Measures	,463 <sup>c</sup>	-,113	,741	1,863	30	30	,047

Two-way mixed effects model where people effects are random and measures effects are fixed.

- The estimator is the same, whether the interaction effect is present or not.
- Type C intraclass correlation coefficients using a consistency definition. The between-measure variance is excluded from the denominator variance.
- This estimate is computed assuming the interaction effect is absent, because it is not estimable otherwise.

### RELIABILITY

```

/VARIABLES=zweitratel_A_Ges Drittrater_A_Ges
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/ICC=MODEL(MIXED) TYPE(CONSISTENCY) CIN=95 TESTVAL=0.

```

## Reliability

### Scale: ALL VARIABLES

#### Case Processing Summary

		N	%
Cases	Valid	31	100,0
	Excluded <sup>a</sup>	0	,0
	Total	31	100,0

- Listwise deletion based on all variables in the procedure.

### Reliability Statistics

Cronbach's Alpha	N of Items
,614	2

**Intraclass Correlation Coefficient**

	Intraclass Correlation <sup>b</sup>	95% Confidence Interval		F Test with True Value 0			
		Lower Bound	Upper Bound	Value	df1	df2	Sig
Single Measures	,443 <sup>a</sup>	,111	,686	2,591	30	30	,006
Average Measures	,614 <sup>c</sup>	,200	,814	2,591	30	30	,006

Two-way mixed effects model where people effects are random and measures effects are fixed.

- a. The estimator is the same, whether the interaction effect is present or not.
- b. Type C intraclass correlation coefficients using a consistency definition. The between-measure variance is excluded from the denominator variance.
- c. This estimate is computed assuming the interaction effect is absent, because it is not estimable otherwise.

GET

FILE='C:\Users\sweyers\MME\Masterarbeit Von der Simulation\Daten\DatenmatrixGesamt H1 und H2.sav'.

DATASET NAME DataSet2 WINDOW=FRONT.