

Attachment 2: Supplementary table

Table S1: Competency level compared by group

Superscript: number of respondents (group with license ≤ 5 yrs. / group with license ≤ 13 yrs.)	I was able to:								I was unable to do it. (highlighted in gray if >50%)	
	teach it to someone else [%]		independently [%]		with guidance when needed [%]		with direct supervision [%]			
	Lic. ≤ 5 yrs.	Lic. ≤ 13 yrs.	Lic. ≤ 5 yrs.	Lic. ≤ 13 yrs.	Lic. ≤ 5 yrs.	Lic. ≤ 13 yrs.	Lic. ≤ 5 yrs.	Lic. ≤ 13 yrs.	Lic. ≤ 5 yrs.	Lic. ≤ 13 yrs.
Examine										
Anal region ^{47/91}	6.4 (n=3)	6.6 (n=6)	29.8 (n=14)	29.7 (n=27)	25.5 (n=12)	26.4 (n=24)	19.1 (n=9)	14.3 (n=13)	19.1 (n=9)	23.1 (n=21)
Extremities ^{47/89}	12.8 (n=6)	7.9 (n=7)	34.0 (n=16)	33.7 (n=30)	29.8 (n=14)	37.1 (n=33)	12.8 (n=6)	10.1 (n=9)	10.6 (n=5)	11.2 (n=10)
Spine ^{46/87}	8.7 (n=4)	5.7 (n=5)	39.1 (n=18)	33.3 (n=29)	23.9 (n=11)	29.9 (n=26)	8.7 (n=4)	10.3 (n=9)	19.6 (n=9)	20.7 (n=18)
Eyes ^{46/88}	/	1.1 (n=1)	15.2 (n=7)	13.6 (n=12)	17.4 (n=8)	13.6 (n=12)	23.9 (n=11)	21.6 (n=19)	43.5 (n=20)	50.0 (n=44)
Skin ^{46/89}	2.2 (n=1)	1.1 (n=1)	17.4 (n=8)	14.6 (n=13)	21.7 (n=10)	25.8 (n=23)	21.7 (n=10)	20.2 (n=18)	37.0 (n=17)	38.2 (n=34)
Nervous system ^{47/89}	10.6 (n=5)	6.7 (n=6)	29.8 (n=14)	27.0 (n=24)	34.0 (n=16)	39.3 (n=35)	17.0 (n=8)	14.6 (n=13)	8.5 (n=4)	12.4 (n=11)
Male genitals including prostate ^{47/90}	2.1 (n=1)	2.2 (n=2)	4.3 (n=2)	4.4 (n=4)	17.0 (n=8)	18.9 (n=17)	12.8 (n=6)	15.6 (n=14)	63.8 (n=30)	58.9 (n=53)
Female breast and lymph nodes ^{46/90}	6.5 (n=3)	3.3 (n=3)	19.6 (n=9)	20.0 (n=18)	26.1 (n=12)	23.3 (n=21)	17.4 (n=8)	23.3 (n=21)	30.4 (n=14)	30.0 (n=27)
Female genitals including speculum ^{45/87}	/	/	2.2 (n=1)	1.1 (n=1)	2.2 (n=1)	1.1 (n=1)	13.3 (n=6)	17.2 (n=15)	82.2 (n=37)	80.5 (n=70)
Neonate ^{46/88}	/	/	6.5 (n=3)	6.8 (n=6)	6.5 (n=3)	9.1 (n=8)	32.6 (n=15)	28.4 (n=25)	54.3 (n=25)	55.7 (n=49)
Infant ^{46/88}	/	/	8.7 (n=4)	4.5 (n=4)	15.2 (n=7)	17.0 (n=15)	19.6 (n=9)	20.5 (n=18)	56.5 (n=26)	58.0 (n=51)
Toddler ^{47/90}	2.1 (n=1)	1.1 (n=1)	6.4 (n=3)	4.4 (n=4)	25.5 (n=12)	23.3 (n=21)	14.9 (n=7)	25.6 (n=23)	51.1 (n=24)	45.6 (n=41)
Sense of balance ^{47/90}	8.5 (n=4)	5.6 (n=5)	31.9 (n=15)	40.0 (n=36)	25.5 (n=12)	22.2 (n=20)	21.3 (n=10)	15.6 (n=14)	12.8 (n=6)	16.7 (n=15)
Geriatric exam & test procedures ^{45/88}	6.7 (n=3)	3.4 (n=3)	11.1 (n=5)	14.8 (n=13)	24.4 (n=11)	20.5 (n=18)	20.0 (n=9)	12.5 (n=11)	37.8 (n=17)	48.9 (n=43)
Perform										
Prick tests ^{43/86}	2.3 (n=1)	1.2 (n=1)	7.0 (n=3)	7.0 (n=6)	7.0 (n=3)	9.3 (n=8)	7.0 (n=3)	11.6 (n=10)	76.7 (n=33)	70.9 (n=61)
Intracutaneous tests ^{43/84}	/	/	2.3 (n=1)	3.6 (n=3)	/	3.6 (n=3)	7.0 (n=3)	6.0 (n=5)	90.7 (n=39)	86.9 (n=73)

Epicutaneous tests ^{43/85}	/	/	4.7 (n=2)	4.7 (n=4)	/	4.7 (n=4)	11.6 (n=5)	7.1 (n=6)	83.7 (n=36)	83.5 (n=71)
Taking samples for pathogen detection ^{46/88}	6.5 (n=3)	6.8 (n=6)	26.1 (n=12)	25.0 (n=22)	19.6 (n=9)	14.8 (n=13)	17.4 (n=8)	17.0 (n=15)	30.4 (n=14)	36.4 (n=32)
Lumbar puncture ^{45/88}	/	/	4.4 (n=2)	2.3 (n=2)	6.7 (n=3)	3.4 (n=3)	8.9 (n=4)	25.0 (n=22)	80.0 (n=36)	69.3 (n=61)
Arterial blood draw ^{46/89}	8.7 (n=4)	7.9 (n=7)	19.6 (n=9)	18.0 (n=16)	10.9 (n=5)	13.5 (n=12)	23.9 (n=11)	20.2 (n=18)	37.0 (n=17)	40.4 (n=36)
Local anesthesia ^{47/89}	6.4 (n=3)	5.6 (n=5)	21.3 (n=10)	16.9 (n=15)	10.6 (n=5)	14.6 (n=13)	12.8 (n=6)	16.9 (n=15)	48.9 (n=23)	46.1 (n=41)
Immobilization techniques/Bandaging ^{46/8}	6.5 (n=3)	8.0 (n=7)	19.6 (n=9)	21.6 (n=19)	17.4 (n=8)	13.6 (n=12)	21.7 (n=10)	21.6 (n=19)	34.8 (n=16)	35.2 (n=31)
Nasal pack ^{46/88}	/	/	4.3 (n=2)	6.8 (n=6)	/	4.5 (n=4)	15.2 (n=7)	12.5 (n=11)	80.4 (n=37)	76.1 (n=67)
Post-mortem examination ^{45/89}	6.7 (n=3)	4.5 (n=4)	13.3 (n=6)	14.6 (n=13)	15.6 (n=7)	20.2 (n=18)	26.7 (n=12)	21.3 (n=19)	37.8 (n=17)	39.3 (n=35)
Insert										
Transurethral bladder catheter ^{46/89}	2.2 (n=1)	6.7 (n=6)	4.3 (n=2)	12.4 (n=11)	17.4 (n=8)	11.2 (n=10)	19.6 (n=9)	19.1 (n=17)	56.5 (n=26)	50.6 (n=45)
Feeding tube ^{44/84}	2.3 (n=1)	4.8 (n=4)	4.5 (n=2)	7.1 (n=6)	6.8 (n=3)	7.1 (n=6)	27.3 (n=12)	26.2 (n=22)	59.1 (n=26)	54.8 (n=46)
Administer medication										
Subcutaneous ^{47/89}	14.9 (n=7)	18.0 (n=16)	31.9 (n=15)	41.6 (n=37)	10.6 (n=5)	5.6 (n=5)	14.9 (n=7)	13.5 (n=12)	27.7 (n=13)	21.3 (n=19)
Intramuscular ^{46/88}	15.2 (n=7)	13.6 (n=12)	28.3 (n=13)	28.4 (n=25)	19.6 (n=9)	19.3 (n=17)	4.3 (n=2)	9.1 (n=8)	32.6 (n=15)	29.5 (n=26)
Intravenous ^{47/91}	17.0 (n=8)	19.8 (n=18)	40.4 (n=19)	46.2 (n=42)	19.1 (n=9)	14.3 (n=13)	8.5 (n=4)	6.6 (n=6)	14.9 (n=7)	13.2 (n=12)
Nasal ^{44/85}	13.6 (n=6)	9.4 (n=8)	31.8 (n=14)	35.3 (n=30)	11.4 (n=5)	8.2 (n=7)	13.6 (n=6)	9.4 (n=8)	29.5 (n=13)	37.6 (n=32)
Canthal ^{45/84}	2.2 (n=1)	1.2 (n=1)	6.7 (n=3)	4.8 (n=4)	2.2 (n=1)	2.4 (n=2)	6.7 (n=3)	4.8 (n=4)	82.2 (n=37)	86.9 (n=73)
Rectal ^{46/90}	10.9 (n=5)	10.0 (n=9)	37.0 (n=17)	43.3 (n=39)	6.5 (n=3)	8.9 (n=8)	10.9 (n=5)	8.9 (n=8)	34.8 (n=16)	28.9 (n=26)
Explain and Demonstrate										
Age-specific peripheral venous catheter ^{47/91}	31.9 (n=15)	26.4 (n=24)	46.8 (n=22)	50.5 (n=46)	2.1 (n=1)	4.4 (n=4)	8.5 (n=4)	6.6 (n=6)	10.6 (n=5)	12.1 (n=11)
Correct use of nebulizer in child ^{47/88}	4.3 (n=2)	4.5 (n=4)	10.6 (n=5)	10.2 (n=9)	10.6 (n=5)	8.0 (n=7)	10.6 (n=5)	8.0 (n=7)	63.8 (n=30)	69.3 (n=61)
Inquire / Describe / Document										
Patients' risk of harm to self and others ^{8/88}	8.7 (n=4)	5.7 (n=5)	23.9 (n=11)	25.0 (n=22)	23.9 (n=11)	29.5 (n=26)	6.5 (n=3)	10.2 (n=9)	37.0 (n=17)	29.5 (n=26)
Psychopathological report ^{46/89}	2.2 (n=1)	1.1 (n=1)	8.7 (n=4)	11.2 (n=10)	37.0 (n=17)	32.6 (n=29)	26.1 (n=12)	20.2 (n=18)	26.1 (n=12)	34.8 (n=31)